Neumotil Vet®

Tilmicosin USP 250 mg/mL





Ensures rapid killing of Mycoplasma & Bacteria effectively

COMPOSITION

Each 100 mL Neumotil vet solution contains- Tilmicosin (as Tilmicosin Phosphate) USP 25 g.

MODE OF ACTION

The antibiotic effects of Neumotil result from the drug binding the 50s ribosome of bacteria and reducing protein synthesis.

INDICATION

Neumotil Vet Oral Solution is indicated for the control and treatment of respiratory infections associated with Neumotil susceptible micro-organisms such as Mycoplasma spp. Pasturella multocida, Actinobacillus, pleuropneumoniae, Actinomyces pyogenes & Mannheimia haemolytica in chickens, calves & turkeys.

DOSAGES & ADMINISTRATION

Poultry: 1mL per 3 liters of drinking water for 3-5 days (75 mg/liter drinking water) Calves: Twice daily, 1 mL per 20 kg body weight via artificial milk for 3 - 5 days. Or as directed by the registered Veterinarian.

CONTRAINDICATION

Contraindicated in poultry hypersensitive to Tilmicosin.

Withdrawal Period

Poultry: Broiler- 12 days, Turkey- 19 days.

STORAGE

Store below 30 ° C, Keep away from light. Keep out of reach of Children.

PACK SIZE

100 mL & 500 mL.