



Rocipro Vet[®]

Ciprofloxacin USP 100 mg/mL

Cure digestive and respiratory diseases effectively

Rocipro Vet[®]

Ciprofloxacin USP 100 mg/mL

100 mL
500 mL
1 Liter

Drug of choice
for diarrhea

An antibiotic with
antioxidant feature

Special Features

A traditional antibiotic
which is highly effective in
respiratory tract & gastrointestinal
tract infections

Significantly reduces
mortality rate in poultry

Comparison of mortality rate among non-treated and Ciprofloxacin treated group




Figure shows that treatment with Ciprofloxacin significantly reduces mortality rate in comparison to non-treated group

■ Mortality Rate-Non Treated Group
■ Mortality Rate-Treated by Ciprofloxacin

Reference: Abdel Ziz, S. A., Abdel Motaal, S. M. A., Abd-Allah, O. E., Sarhan, M. M. I., Pharmacology Dept., Faculty of Vet. Med. Zagazig University, & Animal Health Research Institute, Zagazig Branch. (2016). Concurrent use of ciprofloxacin and metronidazole for controlling of some bacterial infections in broiler chickens. BENHA VETERINARY MEDICAL JOURNAL, 31(2), 83-92. https://bvmlj.journals.ekb.eg/article_31274_ea4fe9f9490df8e93aa3769552cf7c3d.pdf

Indications

- Colibacillosis
- Salmonellosis
- Staphylococcosis
- Pasteurellosis or Fowl cholera
- Coryza
- Mycoplasmosis
- Chronic Respiratory Disease (CRD)

Dosage And Administration

1 mL / 1-2 Liter of drinking water for 3-5 days (8 mg/kg body weight) or as directed by the registered Veterinarian.

Contraindications

Contraindicated to hypersensitivity to Ciprofloxacin or to other quinolones. Chloramphenicol, macrolide and tetracycline should not be used with Ciprofloxacin.

Withdrawal Period

Meat 5 days and Eggs – Not Known

Storage

Store at temperature not exceeding 30°C in a dry place. Protect from light & moisture. Keep out of reach of children.



Manufactured for
Healthcare Pharmaceuticals Ltd.
by NAAFCO Pharma Ltd.
Dhaka, Bangladesh



Marketed by:
Healthcare Formulations Limited
Animal Health Division
Dhaka, Bangladesh