

Neumotil Vet[®]

Tilmicosin USP 250 mg/mL

Ensures rapid killing of
mycoplasma & bacteria
effectively



Special Features

- ✓ Neumotil acts simultaneously as an antibiotic and anti-inflammatory agent
- ✓ Low MIC value range (0.01-0.16 μ g/mL), effective at low concentrations
- ✓ Reaches 4-13 times higher concentrations in air sacs and lungs than other cells, making it highly effective against respiratory tract infections




Neumotil Vet[®]

100 mL
500 mL

Tilmicosin USP 250 mg/mL

Cumulative Feed Conversion Ratio



Infection with Mycoplasma gallisepticum




Figure shows significant reduction in mortality rate of tilmicosin treated group compared to non treated group in Mycoplasma gallisepticum infected broiler chickens

Indication

- Mycoplasmosis
- Chronic Respiratory Disease (CRD)
- Complex Chronic Respiratory Disease (CCRD)
- Pneumonia
- Fowl Cholera (Pasteurellosis)
- Secondary Bacterial Infections

Contraindication

Contraindicated in poultry hypersensitive to Tilmicosin.

Dosages & Administration

1mL per 3 liters of drinking water daily for 3-5 days (15-20mg/kg body weight). Or, as directed by the registered Veterinarian.

Withdrawal Period

Poultry: Broiler-12 days, Turkey-19 days.

website: www.bdhl.com

References:

1. <https://www.researchgate.net/profile/Wafaa-Abd-El-Ghany/publication/45949252>
2. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6358261/>
3. <https://www.sciencedirect.com/science/article/pii/S0022030299753439>



Manufactured for
Healthcare Pharmaceuticals Ltd.
by NAAFCO Pharma Ltd.
Dhaka, Bangladesh



Marketed by
Healthcare Formulations Limited
Animal Health Division
Dhaka, Bangladesh